

Phillips[®] Stool Softener Liquid Gels

Drug Facts	
Active ingredient (in each softgel)	Purpose
Docusate sodium 100 mg.....	Stool softener laxative
Use	
<ul style="list-style-type: none"> • for relief of occasional constipation (irregularity) • this product generally produces a bowel movement in 12 to 72 hours 	
Warnings	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> • stomach pain, nausea, or vomiting • a sudden change in bowel habits that lasts over 14 days 	
Ask a doctor or pharmacist before use if you are presently taking mineral oil	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week 	
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
Directions	
<ul style="list-style-type: none"> • take softgels with a full glass (8 oz) of water 	
adults and children 12 years and older	take 1 to 3 softgels daily or as directed by a doctor. This dose may be taken as a single daily dose or in divided doses.
children 6 to under 12 years	take 1 softgel daily or as directed by a doctor
children under 6 years of age	ask a doctor
Other information	
<ul style="list-style-type: none"> • each softgel contains: sodium 6 mg • very low sodium • store at room temperature. Avoid excessive heat 40°C (104°F). 	
Inactive ingredients FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide	
Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST) or www.bayercare.com	