



Phillips' Milk of Magnesia – Cherry

Drug Facts

Active ingredient (in each 15 mL)	Purpose
Magnesium hydroxide 1200 mg.....	Saline laxative

Use relieves occasional constipation (irregularity). This product usually produces bowel movement in ½ to 6 hours.

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 oz) of liquid with each dose.
- for accurate dosing, only use the dosing cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
---	----------------

children 6 to 11 years	15 mL to 30 mL
------------------------	----------------

children under 6 years	ask a doctor
------------------------	--------------

Other information

- **each 15 mL contains:** magnesium 500 mg, sodium 3 mg
- store at room temperature. Avoid freezing. Close cap tightly after use.

Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C red #28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST)