



Phillips' Concentrated Milk of Magnesia – Strawberry

Drug Facts	
Active ingredient (in each 15 mL)	Purpose
Magnesium hydroxide 2400 mg.....	Saline laxative
Use relieves occasional constipation (irregularity). This product usually produces bowel movement in ½ to 6 hours.	
Warnings	
Do not use if you have ever had an allergic reaction to this product or any of its ingredients	
Ask a doctor before use if you have	
<ul style="list-style-type: none">• kidney disease• a magnesium-restricted diet• stomach pain, nausea, or vomiting• a sudden change in bowel habits that lasts over 14 days	
Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.	
Stop use and ask a doctor if	
<ul style="list-style-type: none">• you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.• you need to use a laxative for more than 1 week	
If pregnant or breast-feeding , ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none">• do not exceed the maximum recommended daily dose in a 24 hour period• shake well before use• dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 oz) of liquid with each dose.• for accurate dosing, only use the dosing cup provided• mL = milliliter• Adults and children 12 years and older: 15 mL to 30 mL• Children under 12 years: ask a doctor	
Other information	
<ul style="list-style-type: none">• each 15 mL contains: magnesium 1000 mg• store at room temperature. Avoid freezing. Close cap tightly after use.	
Inactive ingredients anhydrous citric acid, D&C red #28, flavors, glycerin, propylene glycol, purified water, sodium hypochlorite, sorbitol, sucrose	
Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST)	